arXiv:1301.2139v1 [hep-th] 10 Jan 2013

Triple M-brane configurations and preserved supersymmetries

A. A. Golubtsova$^{1\,,b,c}$ **and V. D. Ivashchuk**$^{2\,,a,b}$

(a) Center for Gravitation and Fundamental Metrology, VNIIMS, 46 Ozyornaya Str., Moscow 119361, Russia

(b) Institute of Gravitation and Cosmology, Peoples' Friendship University of Russia, 6 Miklukho-Maklaya Str., Moscow 117198, Russia

(c) Laboratoire de Univers et Théories (LUTh), Observatoire de Paris, Place Jules Janssen 5, 92190 Meudon, France

Abstract

We investigate all standard triple composite M-brane intersections defined on products of Ricci-flat manifolds for preserving supersymmetries in eleven dimensional $\mathcal{N} = 1$ supergravity. The explicit formulae for computing the numbers of preserved supersymmetries are obtained, which generalize the relations for topologically trivial flat factor spaces presented in the classification by Bergshoeff et al. We obtain certain examples of configurations preserving some fractions of supersymmetries, e.g. containing such factor-spaces as $K3$, $\mathbb{C}^2_*/Z_2$, a four dimensional pp-wave manifold and two-dimensional pseudo-Euclidean manifold $\mathbb{R}^{1,1}_*/Z_2$.

1 Introduction

The developments of [1, 2, 3] have led to a renewed interest in various aspects of supergravity. Classical BPS configurations of intersecting branes play an essential role in studies of non-perturbative superstring and M-theories as well as in establishing and proving new supergravity/gauge correspondences. Non-maximally supersymmetric solutions are important in many applications of superstring dualities. Various intersections of M-branes in 11-dimensional supergravity [4] provide a unified viewpoint, because a large class of solutions describing brane intersections can be obtained by dimensional reduction and duality transformations.

In the basic $M2$- and $M5$-brane solutions, preserving half of supersymetries [5, 6], the worldwolumes have been taken to be flat pseudo-Euclidean spaces $\mathbb{R}^{1,k}$ ($k = 2, 5$) and the transverse spaces have been flat Euclidean ones $\mathbb{R}^r$ ($r = 8, 5$). Nevertheless, the brane configurations defined on the spacetimes with more complicated geometry involved are of interest [7, 8].

First examples of $M2$-brane solutions, partially preserving the supersymmetry, with Ricci-flat 8-dimensional transverse spaces and flat brane worldvolumes $\mathbb{R}^{1,2}$ have been obtained in [9] and [10]. The supersymmetric $M5$-brane solutions with flat transverse spaces

---

[1] siedhe@gmail.com

[2] ivashchuk@mail.ru

$\mathbb{R}^8$ and Ricci-flat 6-dimensional brane wordvolumes have been found in [11], [12] and [13]. In [19] explicit formulae for fractional numbers of supersymmetries for $M2$- and $M5$-brane solutions have been derived. Some new examples have been constructed in [20][1].

In this paper, we study triple orthogonal intersections of composite M-branes defined on the manifold of the form

$$M = M_0 \times M_1 \times \ldots \times M_n, \tag{1.1}$$

where all factor-spaces M_i are Ricci-flat manifolds. It should be noted that the study of the flat case of the factor spaces was undertaken in a variety of works [14]-[18]. In [15] the classification of supersymmetric M-brane configurations on the product manifold with factor spaces $M_i = \mathbb{R}^{k_i}$ was presented. According to this work the amount of preserved supersymmetries is given by

$$\mathcal{N} = 2^{-k} \quad \text{with} \quad k = 1, 2, 3, 4, 5. \tag{1.2}$$

However, the relation (1.2) is not well justified if M-brane configurations are taken into consideration on the product of Ricci-flat manifolds (1.1). In this case the fractional number of supersymmetries $\mathcal{N}$ depends upon several numbers of chiral parallel (i.e. covariantly constant) spinors on certain factor spaces M_i and brane sign factors c_s, which define the orientations of brane worldvolumes.

For clarity, we remind that the metric for orthogonally intersecting M-branes is split into several parts: the common worldvolume, the relative transverese space and the totally transverse one. The classification of possible factor spaces contained in worldvolumes and transverse spaces and admitting parallel spinors can be given in terms of the holonomy groups, see [11],[13],[20] and references therein. While for intersections of two M-branes, as well as for the case of single M2- and M5-branes, one can use the results obtained in works [21],[22], finding non-trivial examples for triple M-brane configurations is complicated by increasing dominance of low-dimensional flat manifolds(of dimensions 1,2 and 3) among factor-spaces.

The purpose of this work is to find out the relations for the amount of preserved supersymmetries for triple M-brane configurations defined on the product of Ricci-flat manifolds. The cases of one and two M-branes were considered in [19] and [20], respectively. Here we deal with non-localized composite brane solutions with zero contributions from Chern-Simons term. However this may be a starting point for future considering of localized brane solutions, which are of interest in view of possible applications by using gravity/gauge correspondence [26].

The structure of the paper is as follows. In section 2 we present a set up, main definitions and notations. Here we use the Propositions 1 and 2 from the previous work [20] which reduce the solutions to generalized Killing equations to a search of parallel (i. e. covariantly constant) spinors on the product manifold (1.1) obeying three algebraic equations. These equations depend upon a brane configuration and brane sign factors. In section 3 we find relations for fractional numbers of preserved supersymmetries for triple M-brane solutions: $M5 \cap M5 \cap M5$, $M2 \cap M2 \cap M5$ and $M2 \cap M5 \cap M5$. For a completeness we start here by considering of three electric branes $M2 \cap M2 \cap M2$ which was performed earlier in [20].

---

[1] In the first two versions of the arxiv publication several examples containing a 2-dimensional flat pseudo-Euclidean submanifold $\mathbb{R}^{1,1}_{*}/Z_2$ with one parallel spinor have been considered.

2 Generalized Killing spinor equations

The bosonic action in 11-dimensional supergravity is given by

$$S_{act} = \int d^{11}z\sqrt{|g|}\left\{R[g] - \frac{1}{2(4!)}F^2\right\} - \frac{1}{6}\int A\wedge F\wedge F, \tag{2.1}$$

where

$$F = dA = \frac{1}{4!}F_{NPQR}dz^N\wedge dz^P\wedge dz^Q\wedge dz^R \tag{2.2}$$

is the 4-form field strength of a 3-form potential A.

The solutions to the equations of motion for the model (2.1) are defined on the (oriented) warped product spin manifold of the form

$$M = M_0\times M_1\times\ldots\times M_n \tag{2.3}$$

with the metric

$$g = e^{2\gamma(x)}g^0 + \sum_{i=1}^{n} e^{2\phi^i(x)}g^i, \tag{2.4}$$

where $g^0 = g^0_{\mu\nu}dx^\mu\otimes dx^\nu$ is a metric on the (oriented) spin manifold M_0 and $g^i = g^i_{m_in_i}dy_i^{m_i}\otimes dy_i^{n_i}$ is a metric on the (oriented) spin manifold M_i, $i = 1,\ldots,n$. We denote $d_\nu = \dim M_\nu$, $\nu = 0,\ldots,n$; $\sum_{\nu=0}^{n} d_\nu = 11$.

The manifold (2.3) allows a frame such that the metric $g = g_{MN}dx^M\otimes dx^N$ ($M,N = 0,\ldots,10$) can be represented in the following form

$$g_{MN} = \eta_{AB}e^A{}_M e^B{}_N, \quad \text{with} \quad \eta_{AB} = \eta^{AB} = \eta_A\delta_{AB}, \tag{2.5}$$

where $e^A = e^A{}_M dx^M$ is the diagonalizing 11-bein, $\eta_A = \pm 1$; $A,B = 0,\ldots,10$. (Here $\eta_A = -1$ only for one value of the index A.)

The required backgrounds must admit 32-component Majorana-spinors ε such that the supersymmetry variation of the gravitino field $\delta\psi_M$ vanishes, i. e.

$$(D_M + B_M)\varepsilon = 0. \tag{2.6}$$

Here

$$D_M = \partial_M + \frac{1}{4}\omega_{ABM}\hat{\Gamma}^A\hat{\Gamma}^B \tag{2.7}$$

is the spinorial covariant derivative, $\hat{\Gamma}^A$ are 32×32 gamma matrices in an orthonormal frame satisfying the Clifford algebra relations

$$\hat{\Gamma}_A\hat{\Gamma}_B + \hat{\Gamma}_B\hat{\Gamma}_A = 2\eta_{AB}\mathbf{1}_{32} \tag{2.8}$$

and $\omega^A{}_{BM}$ is the spin connection and $\omega_{ABM} = \eta_{AC}\omega^C{}_{BM} = -\omega_{BAM}$. (See also the approach of Alekseevsky et al [23]).

In (2.6) B_M is a matrix-valued covector field induced by the 4-form field strength F

$$B_M = \frac{1}{288}\left(\Gamma_M\Gamma^N\Gamma^P\Gamma^Q\Gamma^R - 12\delta^N_M\Gamma^P\Gamma^Q\Gamma^R\right)F_{NPQR}, \tag{2.9}$$

where Γ_M are world gamma matrices obeying

$$\Gamma_M\Gamma_N + \Gamma_M\Gamma_N = 2g_{MN}\mathbf{1}_{32}, \quad \text{with} \quad \Gamma_M = e^A{}_M\hat{\Gamma}_A. \tag{2.10}$$

It should be noted that for any manifold M_l with the metric g^l one considers $k_l \times k_l$ Γ-matrices with $k_l = 2^{[d_l/2]}$ obeying

$$\hat{\Gamma}^{a_l}_{(l)}\hat{\Gamma}^{b_l}_{(l)} + \hat{\Gamma}^{b_l}_{(l)}\hat{\Gamma}^{a_l}_{(l)} = 2\eta^{(l)a_lb_l}\mathbf{1}_{k_l}. \tag{2.11}$$

Here we use alternative double-number notations for indices: $a_j = 1_j, \ldots, (d_j)_j$, where d_j is the dimension of the manifold M_j, $j = 0, \ldots, n$.

The local frame co-vectors are chosen in the following form

$$(e^A{}_M) = \mathrm{diag}\left(e^{\gamma}e^{(0)a}{}_{\mu}, e^{\phi^1}e^{(1)a_1}{}_{m_1}, \ldots, e^{\phi^n}e^{(n)a_n}{}_{m_n}\right), \tag{2.12}$$

where

$$g^0_{\mu\nu} = \eta^{(0)}_{ab}e^{(0)a}{}_{\mu}e^{(0)b}{}_{\nu}, \quad g^i_{m_in_i} = \eta^{(i)}_{a_ib_i}e^{(i)a_i}{}_{m_i}e^{(i)b_i}{}_{n_i}, \tag{2.13}$$

$i = 1, \ldots, n$, and the signature matrix (η_{AB}) in (2.5) can be written down in the components

$$(\eta_{AB}) = \mathrm{diag}\left(\eta^{(0)}_{ab}, \eta^{(1)}_{a_1b_1}, \ldots, \eta^{(n)}_{a_nb_n}\right). \tag{2.14}$$

In (2.13)-(2.14) ($\eta^{(l)}_{a_lb_l}$) is a diagonal signature matrix for the metric g^l, equipped by a (local) frame vectors with components $e^{(l)a_l}{}_{m_l}$, $l = 0, \ldots, n$. We put that

$$\det\left(e^{(l)a_l}{}_{m_l}\right) > 0, \tag{2.15}$$

$l = 0, \ldots, n$, i.e. for any l the oriented set of d_l-beins $e^{(l)a_l}$ has the orientation compatible with the orientation of the manifold M_l.

Henceforth the following notation for the volume d_i-form on the manifolds (M_i, g^i) is used

$$\tau_i \equiv \sqrt{|g^i(y_i)|}dy_i^1 \wedge \ldots \wedge dy_i^{d_i} \tag{2.16}$$

for $i = 1, \ldots, n$. We also denote by $\hat{g}^l$ and $\hat{\tau}_i$ the pull-backs of g^l and τ_i, respectively, to the total manifold M.

In this paper we continue our investigations of composite M-brane solutions [16] (with standard intersection rules) defined on the product of $(n+1)$ Ricci-flat manifolds M_l and governed by several harmonic functions H_s on (M_0, g^0).

It was shown in [20] that the solutions to eqs. (2.6), corresponding to composite M-brane backgrounds, admit a representation in the following form

$$\varepsilon = \left(\prod_{s\in S_e} H_s\right)^{-1/6}\left(\prod_{s\in S_m} H_s\right)^{-1/12}\eta \tag{2.17}$$

where η is a covariantly constant spinor on the spin manifold (2.3) with the metric

$$\bar{g} = \hat{g}^0 + \sum_{i=1}^{n}\hat{g}^i, \tag{2.18}$$

i.e.

$$\bar{D}_M \eta = 0, \tag{2.19}$$

satisfying projection conditions

$$\hat{\Gamma}_{[s]} \eta = c_s \eta, \tag{2.20}$$

$s \in S$, where the brane set S contains two subsets S_e and S_m denoting electric and magnetic brane sets, respectively. Under this convention in (2.17) harmonic functions H_s with $s \in S_e$ correspond to electric $M2$-branes and H_s with $s \in S_m$ correspond to magnetic $M5$-branes.

We remind that in (2.20) $c_s = \pm 1$ are brane sign factors and $\hat{\Gamma}_{[s]}$ are brane operators, $s \in S$, which are defined as follows: for the electric case

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{A_1} \hat{\Gamma}^{A_2} \hat{\Gamma}^{A_3}, \qquad s \in S_e, \tag{2.21}$$

where three indices describe the position of the s-th $M2$-brane worldvolume and for the magnetic case

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{B_1} \hat{\Gamma}^{B_2} \hat{\Gamma}^{B_3} \hat{\Gamma}^{B_4} \hat{\Gamma}^{B_5}, \qquad s \in S_m, \tag{2.22}$$

where five indices B_1, B_2, B_3, B_4, B_5 describe the position not occupied by the s-th $M5$-brane worldvolume. These operators are idempotent, i.e. $(\hat{\Gamma}_{[s]})^2 = \mathbf{1}_{32}$ for all s.

Equations (2.19) are equivalent to the following set of equations [20]

$$\bar{D}^{(l)}_{m_l} \eta = 0, \tag{2.23}$$

$l = 0, \ldots, n$. Here

$$\bar{D}^{(l)}_{m_l} = \partial_{m_l} + \frac{1}{4} \omega^{(l)}_{a_l b_l m_l} \hat{\Gamma}^{a_l} \hat{\Gamma}^{b_l}, \tag{2.24}$$

where $\omega^{(l)}_{a_l b_l m_l} = \eta^{(l)}_{a_l c_l} \omega^{(l) c_l}{}_{b_l m_l}$ and $\omega^{(l) a_l}{}_{b_l m_l}$ are components of the spin connection corresponding to the metric g^l equipped with diagonalizing d_l-bein vectors $e^{(l) a_l}$ and $\hat{\Gamma}^{a_l}_{(l)}$ are gamma matrices corresponding to the manifold M_l and obeying the relations (2.13) $l = 0, ..., n$.

In what follows operators (2.24) will generate the covariant spinorial derivatives corresponding the manifolds M_l

$$D^{(l)}_{m_l} = \partial_{m_l} + \frac{1}{4} \omega^{(l)}_{a_l b_l m_l} \hat{\Gamma}^{a_l}_{(l)} \hat{\Gamma}^{b_l}_{(l)}, \tag{2.25}$$

$l = 0, ..., n$.

In the next section we calculate the fractional numbers $\mathcal{N}$ of unbroken supersymmetries (SUSY) for all triple M-brane configurations. Here we consider generic solutions to the equations (2.6) in the form (2.17). For any configuration we have

$$\mathcal{N} = N/32, \tag{2.26}$$

where N is the dimension of the linear space of solutions to algebraic equations (2.19) and (2.20). (For special non-generic choices of harmonic functions H_s the real fractional numbers $\mathcal{N}$ of unbroken SUSY may be higher than those given by (2.26)).

Remark. In this paper as in [20] we put for simplicity $\varepsilon(z) \in \mathbb{C}^{32}$. The imposing of Majorana condition will give the same number N for the dimension of the real linear space of parallel Majorana spinors obeying (2.19) and (2.20).

3 Triple M-brane configurations

In this section we deduce relations for fractional numbers of supersymmetries preserved by triple M-brane configurations defined on the product manifold (1.1).

3.1 $M2 \cap M2 \cap M2$

Let us consider the configuration of three electric 2-branes intersecting over a point. The configuration which is defined on the manifold

$$M_0 \times M_1 \times M_2 \times M_3 \times M_4, \tag{3.1}$$

with $d_0 = 4$, $d_1 = d_2 = d_3 = 2$ and $d_4 = 1$, can be presented as



Figure 1: $M2 \cap M2 \cap M2$

We define the common worldvolume space by grey, the relative transverse space consists of red, green and blue parts, the totally transverse space is white.

The solution is given by

$$g = H_1^{1/3} H_2^{1/3} H_3^{1/3} \{\hat{g}^0 + H_1^{-1}\hat{g}^1 + H_2^{-1}\hat{g}^2 + H_3^{-1}\hat{g}^3 + H_1^{-1}H_2^{-1}H_3^{-1}\hat{g}^4\}, \tag{3.2}$$

$$F = c_1 dH_1^{-1} \wedge \hat{\tau}_1 \wedge \hat{\tau}_4 + c_2 dH_2^{-1} \wedge \hat{\tau}_2 \wedge \hat{\tau}_4 + c_3 dH_3^{-1} \wedge \hat{\tau}_3 \wedge \hat{\tau}_4, \tag{3.3}$$

where $c_1^2 = c_2^2 = c_3^2 = 1$; H_1, H_2, H_3 are harmonic functions on (M_0, g^0), metrics g^i, $i = 0,1,2,3,4$, are Ricci-flat (the last four metrics are flat). The metrics g^i, $i = 0,1,2,3$, have Euclidean signatures and the metric g^4 has the signature $(-)$. Here we put $M_4 = \mathbb{R}$, $g^4 = -dt \otimes dt$ ($\tau_4 = dt$). The brane sets are $I_1 = \{1,4\}$, $I_2 = \{2,4\}$ and $I_3 = \{3,4\}$.

Using the rules of decomposition for Γ-matrices on product spaces from [25] the set of Γ-matrices can be represented in the following form

$$(\hat{\Gamma}^A) = \begin{array}{ccccccccc} (\hat{\Gamma}^{a_0}_{(0)} & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2 & \otimes & 1, \\ \hat{\Gamma}_{(0)} & \otimes & \hat{\Gamma}^{a_1}_{(1)} & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2 & \otimes & 1, \\ i\hat{\Gamma}_{(0)} & \otimes & \hat{\Gamma}_{(1)} & \otimes & \hat{\Gamma}^{a_2}_{(2)} & \otimes & \mathbf{1}_2 & \otimes & 1, \\ \hat{\Gamma}_{(0)} & \otimes & \hat{\Gamma}_{(1)} & \otimes & \hat{\Gamma}_{(2)} & \otimes & \hat{\Gamma}^{a_3}_{(3)} & \otimes & 1, \\ \hat{\Gamma}_{(0)} & \otimes & \hat{\Gamma}_{(1)} & \otimes & \hat{\Gamma}_{(2)} & \otimes & \hat{\Gamma}_{(3)} & \otimes & 1), \end{array} \tag{3.4}$$

where $\hat{\Gamma}^{a_0}_{(0)}$ are 4×4 gamma matrices corresponding to M_0:

$$\hat{\Gamma}^{a_0}_{(0)}\hat{\Gamma}^{b_0}_{(0)} + \hat{\Gamma}^{b_0}_{(0)}\hat{\Gamma}^{a_0}_{(0)} = 2\delta_{a_0 b_0}\mathbf{1}_4, \tag{3.5}$$

$\hat{\Gamma}^{a_i}_{(i)}$ are 2×2 gamma matrices corresponding to M_i, $i = 1,2,3$:

$$\hat{\Gamma}^{a_i}_{(i)}\hat{\Gamma}^{b_i}_{(i)} + \hat{\Gamma}^{b_i}_{(i)}\hat{\Gamma}^{a_i}_{(i)} = 2\delta_{a_i b_i}\mathbf{1}_2. \tag{3.6}$$

and

$$\hat{\Gamma}_{(0)} = \hat{\Gamma}^{1_0}_{(0)} \dots \hat{\Gamma}^{4_0}_{(0)}, \quad \hat{\Gamma}_{(i)} = \hat{\Gamma}^{1_i}_{(i)} \hat{\Gamma}^{2_i}_{(i)}, \tag{3.7}$$

obey

$$(\hat{\Gamma}_{(0)})^2 = \mathbf{1}_4, \qquad (\hat{\Gamma}_{(i)})^2 = -\mathbf{1}_2. \tag{3.8}$$

The spinor monomial can be written as follows

$$\eta = \eta_0(x) \otimes \eta_1(y_1) \otimes \eta_2(y_2) \otimes \eta_3(y_3) \otimes \eta_4(y_4), \tag{3.9}$$

where $\eta_0 = \eta_0(x)$ is a 4-component spinor on M_0, $\eta_i = \eta_i(y_i)$ is a 2-component spinor on M_i, $i = 1, 2, 3$, and $\eta_4 = \eta_4(y_4)$ is a 1-component spinor on M_4.

Due to (3.4) and (3.8) the covariant derivative acts on the spinor η as follows

$$\begin{aligned} \bar{D}^{(0)}_{m_0}\eta &= (D^{(0)}_{m_0}\eta_0) \otimes \eta_1 \otimes \eta_2 \otimes \eta_3 \otimes \eta_4, & \bar{D}^{(1)}_{m_1}\eta &= \eta_0 \otimes (D^{(1)}_{m_1}\eta_1) \otimes \eta_2 \otimes \eta_3 \otimes \eta_4, \\ \bar{D}^{(2)}_{m_2}\eta &= \eta_0 \otimes \eta_1 \otimes (D^{(2)}_{m_2}\eta_2) \otimes \eta_3 \otimes \eta_4, & \bar{D}^{(3)}_{m_3}\eta &= \eta_0 \otimes \eta_1 \otimes \eta_2 \otimes (D^{(3)}_{m_3}\eta_3) \otimes \eta_4, \\ & & \bar{D}^{(4)}_{m_4}\eta &= \eta_0 \otimes \eta_1 \otimes \eta_2 \otimes \eta_3 \otimes (D^{(4)}_{m_4}\eta_4), \end{aligned} \tag{3.10}$$

where $D^{(i)}_{m_i}$ is a covariant derivative corresponding to M_i, $i = 0, 1, 2, 3$. Here $D^{(4)}_{m_4} = \partial_{m_4}$.

The projection operators corresponding to $M2$-branes are given by

$$\hat{\Gamma}_{[1]} = \hat{\Gamma}^{1_1}\hat{\Gamma}^{2_1}\hat{\Gamma}^{1_4} = -\hat{\Gamma}_{(0)} \otimes \mathbf{1}_2 \otimes \hat{\Gamma}_{(2)} \otimes \hat{\Gamma}_{(3)} \otimes 1 \tag{3.11}$$

for $s = I_1$,

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_2}\hat{\Gamma}^{2_2}\hat{\Gamma}^{1_4} = -\hat{\Gamma}_{(0)} \otimes \hat{\Gamma}_{(1)} \otimes \mathbf{1}_2 \otimes \hat{\Gamma}_{(3)} \otimes 1 \tag{3.12}$$

for $s = I_2$ and

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_3}\hat{\Gamma}^{2_3}\hat{\Gamma}^{1_4} = -\hat{\Gamma}_{(0)} \otimes \hat{\Gamma}_{(1)} \otimes \hat{\Gamma}_{(2)} \otimes \mathbf{1}_2 \otimes 1 \tag{3.13}$$

for $s = I_3$.

The supersymmetry constraints (2.20) are satisfied if

$$\hat{\Gamma}_{(0)}\eta_0 = c_{(0)}\eta_0, \qquad c^2_{(0)} = 1, \tag{3.14}$$

$$\hat{\Gamma}_{(j)}\eta_j = c_{(j)}\eta_j, \qquad c^2_{(j)} = -1, \tag{3.15}$$

$j = 1, 2, 3$, and

$$-c_{(0)}c_{(2)}c_{(3)} = c_1, \qquad -c_{(0)}c_{(1)}c_{(3)} = c_2, \qquad -c_{(0)}c_{(1)}c_{(2)} = c_3. \tag{3.16}$$

Then one obtains the following solution to SUSY equations (2.6) corresponding to the field configuration from (3.2), (3.3)

$$\varepsilon = H_1^{-1/6} H_2^{-1/6} H_3^{-1/6} \eta_0(x) \otimes \eta_1(y_1) \otimes \eta_2(y_2) \otimes \eta_3(y_3) \otimes \eta_4. \tag{3.17}$$

Here η_i, $i = 0, 1, 2, 3$, are chiral parallel spinors defined on M_i, respectively ($D^{(i)}_{m_i}\eta_i = 0$), obeying (3.14), (3.15) and (3.16); η_4 is constant.

Equations (3.16) have the following solutions

$$c_{(0)} = c_1 c_2 c_3, \qquad c_{(j)} = \pm i c_j, \tag{3.18}$$

$j = 1, 2, 3$.

Thus, the number of linear independent solutions given by (3.17)-(3.18) and reads

$$N = 32\mathcal{N} = n_0(c_1c_2c_3)\sum_{c=\pm1} n_1(icc_1)n_2(icc_2)n_3(icc_3), \tag{3.19}$$

where $n_j(c_{(j)})$ is the number of chiral parallel spinors on M_j, $j = 0, 1, 2, 3$; see (3.14) and (3.15).

Examples. Let $M_1 = M_2 = M_3 = \mathbb{R}^2$. The manifold $\mathbb{R}^2$ has one parallel spinor of chirality (+) and one parallel spinor of chirality (-), hence all $n_j(ic) = 1$, $j = 1, 2, 3$, $c = \pm1$, and one gets from (3.19)

$$\mathcal{N} = \frac{1}{16} n_0(c_1c_2c_3). \tag{3.20}$$

It is worth noting that the chirality of the spin η_0 on the manifold M_0 is defined by the product of the brane sign constants c_1, c_2, c_3. For $M_0 = \mathbb{R}^4$ we have $n_0(c) = 2$ and hence $\mathcal{N} = 1/8$ for all values of c_i, $i = 1, 2, 3$.

Consider the case of the curved transverse space. Let $M_0 = K3$, $K3 = CY_2$, which is a 4-dimensional Ricci-flat Kähler manifold with the holonomy group $SU(2) = Sp(1)$. The $K3$ surface has two parallel spinors of the same chirality. We put $n_0(1) = 2$ and $n_0(-1) = 0$. Then we get $\mathcal{N} = 1/8$ if $c_1c_2c_3 = 1$ and $\mathcal{N} = 0$ if $c_1c_2c_3 = -1$. The same result one obtains for the conic space $M_0 = \mathbb{C}^2_*/Z_2$, where $\mathbb{C}^2_* = \mathbb{C}^2 \setminus \{\mathbf{0}\}$. It also has parallel spinors of the same chirality: the pair $(n_0(1), n_0(-1))$ is either $(2, 0)$ or $(0, 2)$ depending on the choice of the spin structure. The completion of $\mathbb{C}^2_*/Z_2$ is the orbifold $\mathbb{C}^2/Z_2$.

3.2 $M5 \cap M5 \cap M5$

According to the classification of M-brane configurations which is presented in [15] there are three possible intersections of three magnetic branes depending on the position of the branes in the bulk space.

(i)

The first case of the solution describing three intersecting $M5$-branes is defined on the manifold of the form

$$M_0 \times M_1 \times M_2 \times M_3 \times M_4, \tag{3.21}$$

where $d_0 = 1$, $d_1 = d_2 = d_3 = 2$, $d_4 = 4$. The configuration is given in Figure 2.



Figure 2: $M5 \cap M5 \cap M5$

The solution reads

$$g = H_1^{2/3} H_2^{2/3} H_3^{2/3} \Big\{ \hat{g}^0 + H_1^{-1}\hat{g}^1 + H_2^{-1}\hat{g}^2 + H_3^{-1}\hat{g}^3 + H_1^{-1}H_2^{-1}H_3^{-1}\hat{g}^4 \Big\}, \tag{3.22}$$

$$F = c_1 (*_0 dH_1) \wedge \hat\tau_2 \wedge \hat\tau_3 + c_2 (*_0 dH_2) \wedge \hat\tau_1 \wedge \hat\tau_3 + c_3 (*_0 dH_3) \wedge \hat\tau_1 \wedge \hat\tau_2, \qquad (3.23)$$

where $c_1^2 = c_2^2 = c_3^2 = 1$; H_1 , H_2 , H_3 are harmonic functions on (M_0, g^0) . The metrics g^i , $i = 0,1,2,3$, have Euclidean signatures and the metric g^4 has the signature $(-,+,+,+)$. The branes sets are $I_1 = \{1,4\}$, $I_2 = \{2,4\}$, $I_3 = \{3,4\}$. Here and in what follows $*_0$ is the Hodge operator on (M_0, g^0) .

Using the rules of decomposition from [25] one can write Γ-matrices in the following form

$$(\hat\Gamma^A) = \begin{array}{ccccccccc} (1 & \otimes & \hat\Gamma_{(1)} & \otimes & \hat\Gamma_{(2)} & \otimes & \hat\Gamma_{(3)} & \otimes & \hat\Gamma_{(4)}, \\ 1 & \otimes & i\hat\Gamma^{a_1}_{(1)} & \otimes & \hat\Gamma_{(2)} & \otimes & \hat\Gamma_{(3)} & \otimes & \hat\Gamma_{(4)}, \\ 1 & \otimes & \mathbf{1}_2 & \otimes & \hat\Gamma^{a_2}_{(2)} & \otimes & \hat\Gamma_{(3)} & \otimes & \hat\Gamma_{(4)}, \\ 1 & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2 & \otimes & i\hat\Gamma^{a_3}_{(3)} & \otimes & \hat\Gamma_{(4)}, \\ 1 & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2 & \otimes & \hat\Gamma^{a_4}_{(4)}) \end{array} \qquad (3.24)$$

Here the operators $\hat\Gamma_{(i)}$, $i = 1,2,3,4$, are given by

$$\hat\Gamma_{(1)} = \hat\Gamma^{1_1}_{(1)}\hat\Gamma^{2_1}_{(1)}, \quad \hat\Gamma_{(2)} = \hat\Gamma^{1_2}_{(2)}\hat\Gamma^{2_2}_{(2)}, \quad \hat\Gamma_{(3)} = \hat\Gamma^{1_3}_{(3)}\hat\Gamma^{2_3}_{(3)}, \quad \hat\Gamma_{(4)} = \hat\Gamma^{1_4}_{(4)}\hat\Gamma^{2_4}_{(4)}\hat\Gamma^{3_4}_{(4)}\hat\Gamma^{4_4}_{(4)} \qquad (3.25)$$

obey

$$(\hat\Gamma_{(i)})^2 = -\mathbf{1}_2, \quad (\hat\Gamma_{(4)})^2 = -\mathbf{1}_4 \qquad (3.26)$$

with $i = 1,2,3$. $\hat\Gamma^{a_1}_{(1)}, \hat\Gamma^{a_2}_{(2)}, \hat\Gamma^{a_3}_{(3)}$ are 2×2 Γ-matrices corresponding to M_1 , M_2 , M_3 respectively, $\hat\Gamma^{a_4}_{(4)}$ is a set of gamma matrices corresponding to M_4 .

The covariant derivatives can be written down as

$$\begin{aligned} \bar D^{(1)}_{m_1} &= \partial_{m_1} + \frac{1}{4}\omega^{(1)}_{a_1 b_1 m_1}\left(1 \otimes \hat\Gamma^{a_1}_{(1)}\hat\Gamma^{b_1}_{(1)} \otimes \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \mathbf{1}_4\right), \\ \bar D^{(2)}_{m_2} &= \partial_{m_2} + \frac{1}{4}\omega^{(2)}_{a_2 b_2 m_2}\left(1 \otimes \mathbf{1}_2 \otimes \hat\Gamma^{a_2}_{(2)}\hat\Gamma^{b_2}_{(2)} \otimes \mathbf{1}_2 \otimes \mathbf{1}_4\right), \\ \bar D^{(3)}_{m_3} &= \partial_{m_3} + \frac{1}{4}\omega^{(3)}_{a_3 b_3 m_3}\left(1 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \hat\Gamma^{a_3}_{(3)}\hat\Gamma^{b_3}_{(3)} \otimes \mathbf{1}_4\right), \\ \bar D^{(4)}_{m_4} &= \partial_{m_4} + \frac{1}{4}\omega^{(4)}_{a_4 b_4 m_4}\left(1 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \hat\Gamma^{a_4}_{(4)}\hat\Gamma^{b_4}_{(4)}\right), \end{aligned} \qquad (3.27)$$

where $\omega^{(i)a_i}{}_{b_i c_i}$ are components of the spin connection corresponding to the manifold M_i , $D^{(i)}_{m_i}$ are covariant derivatives corresponding to M_i , $i = 1,2,3,4$, $\bar D^{(0)}_{m_0} = \partial_{m_0}$ and $D^{(0)}_{m_0} = \partial_{m_0}$.

Let η be represented in the following form

$$\eta = \eta_0(x) \otimes \eta_1(y_1) \otimes \eta_2(y_2) \otimes \eta_3(y_3) \otimes \eta_4(y_4), \qquad (3.28)$$

where $\eta_0(x)$ is a 1-component spinor on M_0 , $\eta_i = \eta_i(y_i)$ is a 2-component spinor on M_i , $i = 1,2,3$, $\eta_4 = \eta_4(y_4)$ is a 4-component spinor on M_4 .

Then for spinorial covariant derivatives we get relations (3.10).

The operators corresponding to the $M5$-branes read

$$\hat\Gamma_{[s]} = \hat\Gamma^{1_0}\hat\Gamma^{1_2}\hat\Gamma^{2_2}\hat\Gamma^{1_3}\hat\Gamma^{2_3} = 1 \otimes \hat\Gamma_{(1)} \otimes \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \hat\Gamma_{(4)} \qquad (3.29)$$

for $s = I_1$,

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_0}\hat{\Gamma}^{1_1}\hat{\Gamma}^{2_1}\hat{\Gamma}^{1_3}\hat{\Gamma}^{2_3} = 1 \otimes \mathbf{1}_2 \otimes \hat{\Gamma}_{(2)} \otimes \mathbf{1}_2 \otimes \hat{\Gamma}_{(4)} \tag{3.30}$$

for $s = I_2$ and

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_0}\hat{\Gamma}^{1_1}\hat{\Gamma}^{2_1}\hat{\Gamma}^{1_2}\hat{\Gamma}^{2_2} = 1 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \hat{\Gamma}_{(3)} \otimes \hat{\Gamma}_{(4)} \tag{3.31}$$

for $s = I_3$.

Thus the spinor η obeys the projection conditions (2.20) if

$$\hat{\Gamma}_{(j)}\eta_j = c_{(j)}\eta_j, \qquad c_{(j)}^2 = -1, \tag{3.32}$$

$j = 1, 2, 3, 4$, and

$$c_{(1)}c_{(4)} = c_1, \quad c_{(2)}c_{(4)} = c_2, \quad c_{(3)}c_{(4)} = c_3. \tag{3.33}$$

The solution to the generalized Killing equations corresponding to the field configuration from (3.22), (3.23) can be represented in the following form

$$\varepsilon = \left(\prod_{s=1}^{3} H_s^{-\frac{1}{12}}\right) \eta_0 \otimes \eta_1(y_1) \otimes \eta_2(y_2) \otimes \eta_2(y_3) \otimes \eta_4(y_4), \tag{3.34}$$

where η_i, $i = 1, 2, 3, 4$, are parallel spinors defined on M_i, respectively $\left(D_{m_i}^{(i)}\eta_i = 0\right)$, obeying (3.32) and (3.33), and η_0 is a constant one-dimensional spinor on M_0.

Eqs. (3.33) have the following solutions

$$c_{(1)} = -ic_1, \quad c_{(2)} = -ic_2, \quad c_{(3)} = -ic_3, \quad c_{(4)} = i, \tag{3.35}$$

or

$$c_{(1)} = ic_1, \quad c_{(2)} = ic_2, \quad c_{(3)} = ic_3, \quad c_{(4)} = -i. \tag{3.36}$$

The number of linear independent solutions given by (3.34), (3.35) and (3.36) reads

$$N = 32\mathcal{N} = n_1(-ic_1)n_2(-ic_2)n_3(-ic_3)n_4(i) + n_1(ic_1)n_2(ic_2)n_3(ic_3)n_4(-i), \tag{3.37}$$

where $n_j(c_j)$ is the number of chiral parallel spinors on M_j, $j = 1, 2, 3, 4$.

Examples. Let $M_0 = \mathbb{R}$ and $M_1 = M_2 = M_3 = \mathbb{R}^2$. Then all $n_j(c) = 1$, $j = 1, 2, 3$, with $c = \pm i$, and hence we get from (3.37)

$$N = 32\mathcal{N} = n_4(i) + n_4(-i). \tag{3.38}$$

For the case when the common worldvolume is coinciding with Minkowski space $M_4 = \mathbb{R}^{1,3}$, we have $n_4(c) = 2$ and hence $\mathcal{N} = 1/8$ for all values of c_i, $i = 1, 2, 3$.

If the $M5$-branes intersect by the manifold $M_4 = (\mathbb{R}^{1,1}_*/Z_2) \times \mathbb{R}^2$, where $\mathbb{R}^{1,1}_* = \mathbb{R}^{1,1} \setminus \{\mathbf{0}\}$. Since $\mathbb{R}^{1,1}_*$ has only one parallel spinor (left or right) depending of the choice of the spin structure, one obtains $n_4(c) = 1$ for any c and hence $\mathcal{N} = 1/16$.

The same result takes place when (M_4, g^4) is a 4-dimensional Ricci-flat pp-wave space from [13] with the holonomy group $H = \mathbb{R}^2$ (see [24]). In this case $(n_4(i), n_4(-i)) = (1, 1)$ and $\mathcal{N} = 1/16$.

(ii)



Figure 3: $M5 \cap M5 \cap M5$

The second possible configuration of three $M5$-branes is pairwise intersection over 3-branes defined on the manifold of the form

$$M_0 \times M_1 \times M_2 \times M_3 \times M_4, \tag{3.39}$$

where $d_0 = 3$, $d_1 = d_2 = d_3 = d_4 = 2$. In Figure 3 one presents the intersection of three magnetic branes.

The metric and the 4-form field strength can be represented in the following form

$$g = H_1^{2/3} H_2^{2/3} H_3^{2/3} \Big\{ \hat{g}^0 + H_2^{-1} H_3^{-1} \hat{g}^1 + H_1^{-1} H_3^{-1} \hat{g}^2 + H_1^{-1} H_2^{-1} \hat{g}^3 + H_1^{-1} H_2^{-1} H_3^{-1} \hat{g}^4 \Big\}, \tag{3.40}$$

$$F = c_1 (*_0 dH_1) \wedge \hat{\tau}_1 + c_2 (*_0 dH_2) \wedge \hat{\tau}_2 + c_3 (*_0 dH_3) \wedge \hat{\tau}_3, \tag{3.41}$$

where $c_1^2 = c_2^2 = c_3^2 = 1$; H_1, H_2, H_3 are harmonic functions on (M_0, g^0). The metrics g^i, $i = 0,1,2,3$, have Euclidean signatures and the metric g^4 has the signature $(-,+)$. The branes sets are $I_1 = \{2,3,4\}$, $I_2 = \{1,3,4\}$, $I_3 = \{1,2,4\}$.

The gamma matrices can be split in the following form

$$(\hat{\Gamma}^A) = \begin{array}{ccccccccc} (i\hat{\Gamma}^{a_0}_{(0)} & \otimes & \hat{\Gamma}_{(1)} & \otimes & \hat{\Gamma}_{(2)} & \otimes & \hat{\Gamma}_{(3)} & \otimes & \hat{\Gamma}_{(4)}, \\ \mathbf{1}_2 & \otimes & \hat{\Gamma}^{a_1}_{(1)} & \otimes & \hat{\Gamma}_{(2)} & \otimes & \hat{\Gamma}_{(3)} & \otimes & \hat{\Gamma}_{(4)}, \\ \mathbf{1}_2 & \otimes & \mathbf{1}_2 & \otimes & i\hat{\Gamma}^{a_2}_{(2)} & \otimes & \hat{\Gamma}_{(3)} & \otimes & \hat{\Gamma}_{(4)}, \\ \mathbf{1}_2 & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2 & \otimes & \hat{\Gamma}^{a_3}_{(3)} & \otimes & \hat{\Gamma}_{(4)}, \\ \mathbf{1}_2 & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2 & \otimes & \hat{\Gamma}^{a_4}_{(4)}), \end{array} \tag{3.42}$$

where

$$\hat{\Gamma}_{(0)} = \hat{\Gamma}^{1_0}_{(0)} \hat{\Gamma}^{2_0}_{(0)} \hat{\Gamma}^{3_0}_{(0)}, \quad \hat{\Gamma}_{(1)} = \hat{\Gamma}^{1_1}_{(1)} \hat{\Gamma}^{2_1}_{(1)} \quad \hat{\Gamma}_{(2)} = \hat{\Gamma}^{1_2}_{(2)} \hat{\Gamma}^{2_2}_{(2)},$$
$$\hat{\Gamma}_{(3)} = \hat{\Gamma}^{1_3}_{(3)} \hat{\Gamma}^{2_3}_{(3)}, \quad \hat{\Gamma}_{(4)} = \hat{\Gamma}^{1_4}_{(4)} \hat{\Gamma}^{2_4}_{(4)}. \tag{3.43}$$

satisfy

$$(\hat{\Gamma}_{(0)})^2 = (\hat{\Gamma}_{(1)})^2 = (\hat{\Gamma}_{(2)})^2 = (\hat{\Gamma}_{(3)})^2 = -\mathbf{1}_2, \quad (\hat{\Gamma}_{(4)})^2 = \mathbf{1}_2. \tag{3.44}$$

Here $\hat{\Gamma}^{a_i}_{(i)}$ are 2×2 gamma matrices corresponding to M_i, $i = 0,1,2,3,4$. One can write down the gamma matrices corresponding to M_0 as $\left(\hat{\Gamma}^{a_0}_{(0)}\right) = (\sigma_1, \sigma_2, \sigma_3)$ and hence

$$\hat{\Gamma}_{(0)} = i\mathbf{1}_2. \tag{3.45}$$

Here we put

$$\eta = \eta_0(x) \otimes \eta_1(y_1) \otimes \eta_2(y_2) \otimes \eta_3(y_3) \otimes \eta_4(y_4), \tag{3.46}$$

where $\eta_0 = \eta_0(x)$ is a 2-component spinor on M_0, $\eta_i = \eta_i(y_i)$ is a 2-component spinor on M_i, $i = 1,2,3,4$.

The factorization relations (3.10) are valid for spinorial covariant derivatives where

$$\bar{D}^{(0)}_{m_0} = \partial_{m_0} + \frac{1}{4}\omega^{(0)}_{a_0 b_0 m_0}\left(\hat{\Gamma}^{a_0}_{(0)}\hat{\Gamma}^{b_0}_{(0)} \otimes \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2\right),$$

$$\ldots$$

$$\bar{D}^{(4)}_{m_4} = \partial_{m_4} + \frac{1}{4}\omega^{(4)}_{a_4 b_4 m_4}\left(\mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \hat{\Gamma}^{a_4}_{(4)}\hat{\Gamma}^{b_4}_{(4)}\right), \qquad (3.47)$$

$D^{(i)}_{m_i}$ are covariant derivatives corresponding to M_i, $i = 0,1,2,3,4$.

Due to (3.42) and (3.43)-(3.45) the brane operators corresponding to the magnetic branes write down as

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_0}\hat{\Gamma}^{2_0}\hat{\Gamma}^{3_0}\hat{\Gamma}^{1_1}\hat{\Gamma}^{2_1} = \mathbf{1}_2 \otimes \mathbf{1}_{(2)} \otimes \hat{\Gamma}_{(2)} \otimes \hat{\Gamma}_{(3)} \otimes \hat{\Gamma}_{(4)} \qquad (3.48)$$

for $s = I_1$,

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_0}\hat{\Gamma}^{2_0}\hat{\Gamma}^{3_0}\hat{\Gamma}^{1_2}\hat{\Gamma}^{2_2} = \mathbf{1}_2 \otimes \hat{\Gamma}_{(1)} \otimes \mathbf{1}_2 \otimes \hat{\Gamma}_{(3)} \otimes \hat{\Gamma}_{(4)} \qquad (3.49)$$

for $s = I_2$,

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_0}\hat{\Gamma}^{2_0}\hat{\Gamma}^{3_0}\hat{\Gamma}^{1_3}\hat{\Gamma}^{2_3} = \mathbf{1}_2 \otimes \hat{\Gamma}_{(1)} \otimes \hat{\Gamma}_{(2)} \otimes \mathbf{1}_2 \otimes \hat{\Gamma}_{(4)} \qquad (3.50)$$

for $s = I_3$.

The supersymmetry restrictions (2.20) are satisfied if

$$\hat{\Gamma}_{(1)}\eta_1 = c_1\eta_1, \qquad c^2_{(1)} = 1, \qquad (3.51)$$

$$\hat{\Gamma}_{(j)}\eta_j = c_{(j)}\eta_j, \qquad c^2_{(j)} = -1, \qquad (3.52)$$

with $j = 0,2,3,4$. Then the conditions for the chirality constants are given by

$$c_{(1)}c_{(3)}c_{(4)} = c_1, \quad c_{(1)}c_{(2)}c_{(3)} = c_2, \quad c_{(1)}c_{(2)}c_{(4)} = c_3. \qquad (3.53)$$

We get the following solution to equations (2.6) corresponding to the field configuration from (3.40)-(3.41)

$$\varepsilon = H_1^{-1/12}H_2^{-1/12}H_3^{-1/12}\eta_0(x) \otimes \eta_1(y_1) \otimes \eta_2(y_2) \otimes \eta_2(y_3) \otimes \eta_4(y_4). \qquad (3.54)$$

Here η_i, $i = 0,1,2,3,4$, are chiral parallel spinors defined on M_i $\left(D^{(i)}_{m_i}\eta_i = 0\right)$, obeying (3.51)-(3.53). Equations (3.53) have the following solutions

$$c_{(1)} = \pm i c_2 c_3, \quad c_{(2)} = \pm i c_1 c_2, \quad c_{(3)} = \pm i, \quad c_{(4)} = -c_1 c_2 c_3. \qquad (3.55)$$

The number of linear independent solutions is

$$N = 32\mathcal{N} = n_0 n_4(-c_1c_2c_3)\sum_{c=\pm 1} n_1(icc_2c_3)n_2(icc_1c_2)n_3(ic), \qquad (3.56)$$

where $n_j(c_j)$ is the number of chiral parallel spinors on M_j, $j = 1,2,3,4$, n_0 is the number of parallel spinors on M_0.

Examples. Consider the case when all factor spaces are flat: $M_0 = \mathbb{R}^3$, $M_1 = M_2 = M_3 = \mathbb{R}^2$. Then all $n_j(ic) = 1$, $j = 1,2,3$, $c = \pm 1$. The amount of preserved supersymmetries is given by

$$\mathcal{N} = \frac{1}{8} n_4(-c_1 c_2 c_3). \tag{3.57}$$

Let $M_4 = \mathbb{R}^{1,1}$, then we obtain $\mathcal{N} = \dfrac{1}{8}$ for any choice of brane sign factors. If $M_4 = \mathbb{R}^{1,1}_*/Z_2$ with $n_4(1) = 1$ and $n_4(-1) = 0$ we get $\mathcal{N} = \dfrac{1}{8}$ for $c_1 c_2 c_3 = -1$ and $\mathcal{N} = 0$ otherwise.

(iii)

The third intersection of magnetic branes is defined on the manifold

$$M_0 \times M_1 \times M_2 \times M_3 \times M_4 \times M_5 \times M_6 \times M_7, \tag{3.58}$$

where $d_0 = 2$, $d_1 = d_2 = d_3 = d_4 = d_5 = d_6 = 1$, $d_7 = 3$. For this configuration we have the following picture



Figure 4: $M5 \cap M5 \cap M5$

The metric of three intersecting $M5$ branes reads now

$$g = H_1^{2/3} H_2^{2/3} H_3^{2/3} \Big\{ \hat{g}^0 + H_1^{-1}\hat{g}^1 + H_2^{-1}\hat{g}^2 + H_3^{-1}\hat{g}^3 + H_2^{-1}H_3^{-1}\hat{g}^4 + H_1^{-1}H_3^{-1}\hat{g}^5 + H_1^{-1}H_2^{-1}\hat{g}^6 + H_1^{-1}H_2^{-1}H_3^{-1}\hat{g}^7 \Big\}. \tag{3.59}$$

The corresponding field strength is

$$F = c_1(*_0 dH_1) \wedge \hat{\tau}_2 \wedge \hat{\tau}_3 \wedge \hat{\tau}_4 + c_2(*_0 dH_2) \wedge \hat{\tau}_1 \wedge \hat{\tau}_3 \wedge \hat{\tau}_5 + c_3(*_0 dH_3) \wedge \hat{\tau}_1 \wedge \hat{\tau}_2 \wedge \hat{\tau}_6, \tag{3.60}$$

where $c_1^2 = c_2^2 = c_3^2 = 1$; H_1, H_2, H_3 are harmonic functions on (M_0, g^0). The metrics g^i, $i = 0,1,2,3,4,5,6$, have Euclidean signatures and the metric g^7 has the signature $(-,+,+)$. The branes sets are $I_1 = \{1,5,6,7\}$, $I_2 = \{2,4,6,7\}$, $I_3 = \{3,4,5,7\}$.

Under the decomposition rules the set of gamma matrices can be presented in the form

$$(\hat{\Gamma}^A) = \begin{array}{lllllllllllllllllllll}
(\hat{\Gamma}^{a_0}_{(0)} & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2, \\
i\hat{\Gamma}_{(0)} & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & \hat{\Gamma}_{(7)} & \otimes & \sigma_3 & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2, \\
i\hat{\Gamma}_{(0)} & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & \hat{\Gamma}_{(7)} & \otimes & \sigma_1 & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2, \\
i\hat{\Gamma}_{(0)} & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & \hat{\Gamma}_{(7)} & \otimes & \sigma_2 & \otimes & \sigma_3 & \otimes & \mathbf{1}_2, \\
i\hat{\Gamma}_{(0)} & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & \hat{\Gamma}_{(7)} & \otimes & \sigma_2 & \otimes & \sigma_1 & \otimes & \mathbf{1}_2, \\
i\hat{\Gamma}_{(0)} & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & \hat{\Gamma}_{(7)} & \otimes & \sigma_2 & \otimes & \sigma_2 & \otimes & \sigma_3, \\
i\hat{\Gamma}_{(0)} & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & \hat{\Gamma}_{(7)} & \otimes & \sigma_2 & \otimes & \sigma_2 & \otimes & \sigma_1, \\
i\hat{\Gamma}_{(0)} & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & 1 & \otimes & \hat{\Gamma}^{a_7}_{(7)} & \otimes & \sigma_2 & \otimes & \sigma_2 & \otimes & \sigma_2).
\end{array} \tag{3.61}$$

Here the operators

$$\hat{\Gamma}_{(0)} = \hat{\Gamma}^{1_0}_{(0)} \hat{\Gamma}^{2_0}_{(0)}, \quad \hat{\Gamma}_{(7)} = \hat{\Gamma}^{1_7}_{(7)} \hat{\Gamma}^{2_7}_{(7)} \hat{\Gamma}^{3_7}_{(7)} \tag{3.62}$$

obey

$$(\hat{\Gamma}_{(0)})^2 = -\mathbf{1}_2, \quad (\hat{\Gamma}_{(7)})^2 = \mathbf{1}_2. \tag{3.63}$$

The gamma matrices corresponding to M_0 and M_7 manifolds can be written down in the form $\left(\hat{\Gamma}^{a_0}_{(0)}\right) = (\sigma_1, \sigma_2)$, $\hat{\Gamma}_{(0)} = i\sigma_3$, $\left(\hat{\Gamma}^{a_7}_{(7)}\right) = (i\sigma_1, \sigma_2, \sigma_3)$, $\hat{\Gamma}_{(7)} = -\mathbf{1}_2$, respectively.

We put the following relation for the 32-component spinor

$$\eta = \eta_0(x) \otimes \eta_1(y_1) \otimes \eta_2(y_2) \otimes \eta_3(y_3) \otimes \eta_4(y_4) \otimes \eta_5(y_5) \otimes \eta_6(y_6) \otimes \eta_7(y_7) \otimes \chi, \tag{3.64}$$

where $\eta_i = \eta_i(y_i)$ is a 1-component spinor on M_i , $i = 1, 2, 3, 4, 5, 6$, $\eta_0 = \eta_0(x)$ is a 2-component spinor on M_0 , $\eta_7 = \eta_7(y_7)$ is a 2-component spinor on M_7 and

$$\chi = \sum_{r=1}^{8} \chi_1^r \otimes \chi_2^r \otimes \chi_3^r \tag{3.65}$$

belongs to $V = \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$; $\chi_1^r, \chi_2^r, \chi_3^r$ are “auxiliary” 2-dimensional spinors.

Here the covariant derivatives act on η as

$$\begin{aligned} \bar{D}^{(0)}_{m_0} \eta &= \left(D^{(0)}_{m_0} \eta_0\right) \otimes \eta_1 \otimes \eta_2 \otimes \eta_3 \otimes \eta_4 \otimes \eta_5 \otimes \eta_6 \otimes \eta_7 \otimes \chi, \\ & \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \ldots \\ \bar{D}^{(7)}_{m_7} \eta &= \eta_0 \otimes \eta_1 \otimes \eta_2 \otimes \eta_3 \otimes \eta_4 \otimes \eta_5 \otimes \eta_6 \otimes \left(D^{(7)}_{m_7} \eta_7\right) \otimes \chi, \end{aligned} \tag{3.66}$$

where

$$\begin{aligned} \bar{D}^{(0)}_{m_0} &= \partial_{m_0} + \frac{1}{4} w^{(0)}_{a_0 b_0 m_0} \left(\hat{\Gamma}^{a_0}_{(0)} \hat{\Gamma}^{b_0}_{(0)} \otimes 1 \otimes 1 \otimes 1 \otimes 1 \otimes 1 \otimes 1 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2\right), \\ \bar{D}^{(7)}_{m_7} &= \partial_{m_7} + \frac{1}{4} w^{(7)}_{a_7 b_7 m_7} \left(\mathbf{1}_2 \otimes 1 \otimes 1 \otimes 1 \otimes 1 \otimes 1 \otimes 1 \otimes \hat{\Gamma}^{a_7}_{(7)} \hat{\Gamma}^{b_7}_{(7)} \otimes \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2\right), \end{aligned} \tag{3.67}$$

$\bar{D}^{(i)}_{m_i} = D^{(i)}_{m_i} = \partial_{m_i}$, $i = 1, 2, 3, 4, 5, 6$. $D^{(i)}_{m_i}$ are covariant derivatives corresponding to M_i , $i = 0, 7$. Using (3.61) one can write down the projectors for $M5$-branes as

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_0} \hat{\Gamma}^{2_0} \hat{\Gamma}^{1_2} \hat{\Gamma}^{1_3} \hat{\Gamma}^{1_4} = -\mathbf{1}_2 \otimes 1 \otimes 1 \otimes 1 \otimes 1 \otimes 1 \otimes 1 \otimes \hat{\Gamma}_{(7)} \otimes B_1 \tag{3.68}$$

for $s = I_1$,

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_0} \hat{\Gamma}^{2_0} \hat{\Gamma}^{1_1} \hat{\Gamma}^{1_3} \hat{\Gamma}^{1_5} = -\mathbf{1}_2 \otimes 1 \otimes 1 \otimes 1 \otimes 1 \otimes 1 \otimes 1 \otimes \hat{\Gamma}_{(7)} \otimes B_2 \tag{3.69}$$

for $s = I_2$,

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_0} \hat{\Gamma}^{2_0} \hat{\Gamma}^{1_1} \hat{\Gamma}^{1_2} \hat{\Gamma}^{1_6} = -\mathbf{1}_2 \otimes 1 \otimes 1 \otimes 1 \otimes 1 \otimes 1 \otimes 1 \otimes \hat{\Gamma}_{(7)} \otimes B_3 \tag{3.70}$$

for $s = I_3$.

Here we denote by B_s the following self-adjoint commuting idempotent (i.e. $B_s^2 = \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2$) operators acting on the 8-dimensional Hilbert space $V = \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$

$$B_1 = -\sigma_1 \otimes \sigma_2 \otimes \mathbf{1}_2, \quad B_2 = \sigma_3 \otimes \sigma_1 \otimes \sigma_3, \quad B_3 = -\mathbf{1}_2 \otimes \sigma_2 \otimes \sigma_1. \tag{3.71}$$

Due to Proposition from Appendix there exists a basis of eigenvectors $\psi_{\varepsilon_1,\varepsilon_2,\varepsilon_3}$ in V with $\varepsilon_1 = \pm 1, \varepsilon_2 = \pm 1, \varepsilon_3 = \pm 1$, obeying

$$B_s \psi_{\varepsilon_1,\varepsilon_2,\varepsilon_3} = \varepsilon_s \psi_{\varepsilon_1,\varepsilon_2,\varepsilon_3}, \tag{3.72}$$

$s = 1, 2, 3$.

Let the gamma matrices corresponding to the 3-dimensional manifold M_7 be chosen as follows $\hat{\Gamma}^{a_7}_{(7)} = (i\sigma_1, \sigma_2, \sigma_3)$, and hence $\hat{\Gamma}_{(7)} = -\mathbf{1}_2$.

Then the solutions to SUSY equations (2.6) corresponding to the field configuration from (3.59), (3.60) are generated by the following set of monomial solutions

$$\varepsilon = H_1^{-1/12} H_2^{-1/12} H_3^{-1/12} \eta_0(x) \otimes \eta_1 \otimes \eta_2 \otimes \eta_3 \otimes \eta_4 \otimes \eta_5 \otimes \eta_6 \otimes \eta_7(y_7) \otimes \psi_{\varepsilon_1,\varepsilon_2,\varepsilon_3}, \tag{3.73}$$

where $\eta_0(x)$ and $\eta_7(y_7)$ are parallel spinors defined on M_0 and M_7, respectively, η_i are constant 1-dimensional spinors, $i = 1, 2, 3, 4, 5, 6$. Here ε_s parameters obey the relations

$$\varepsilon_s = c_s, \tag{3.74}$$

$s = 1, 2, 3$, following from relations (2.20), (3.65) with $\chi = \psi_{\varepsilon_1,\varepsilon_2,\varepsilon_3}$, (3.68), (3.69), (3.70) and (3.72).

The number of linear independent solutions given by (3.73) and (3.74) is

$$N = 32\mathcal{N} = n_0 n_7, \tag{3.75}$$

where n_j is the number of parallel spinors on M_j, $j = 0, 7$.

Examples. For $M_0 = \mathbb{R}^2$ and $M_7 = \mathbb{R}^{1,2}$ we get from (3.19) $\mathcal{N} = 1/8$ in agreement with [15].

If we put $M_7 = (\mathbb{R}^{1,1}_*/Z_2) \times \mathbb{R}$ instead of 3-dimensional analogue of Minkowski space $M_7 = \mathbb{R}^{1,2}$ one obtains $\mathcal{N} = 1/16$. This result does not depend upon the brane sign factors.

3.3 $M2 \cap M2 \cap M5$

Let us consider the composite configuration of two electric branes each intersecting $M5$-brane over a string with the two strings intersecting over a point. $M2 \cap M2 \cap M5$-solution is defined on the manifold

$$M_0 \times M_1 \times M_2 \times M_3 \times M_4 \times M_5 \times M_6, \tag{3.76}$$

where $d_0 = 3$, $d_1 = d_2 = d_4 = d_5 = d_6 = 1$ and $d_3 = 3$.

The intersection is given in Figure 5.



Figure 5: $M2 \cap M2 \cap M5$

The metric and the 4-form field strength corresponding to the intersection of two $M2$-branes and one $M5$-brane can be represented in the following form

$$g = H_1^{1/3} H_2^{1/3} H_3^{2/3} \Big\{ \hat{g}^0 + H_1^{-1} \hat{g}^1 + H_2^{-1} \hat{g}^2 + H_3^{-1} \hat{g}^3 + H_2^{-1} H_3^{-1} \hat{g}^4 + \\ H_1^{-1} H_3^{-1} \hat{g}^5 + H_1^{-1} H_2^{-1} H_3^{-1} \hat{g}^6 \Big\}, \qquad (3.77)$$

$$F = c_1 dH_1^{-1} \wedge \hat{\tau}_1 \wedge \hat{\tau}_5 \wedge \hat{\tau}_6 + c_2 dH_2^{-1} \wedge \hat{\tau}_2 \wedge \hat{\tau}_4 \wedge \hat{\tau}_6 + c_3 (*_0 dH_3) \wedge \hat{\tau}_1 \wedge \hat{\tau}_2, \qquad (3.78)$$

where $c_1^2 = c_2^2 = c_3^2 = 1$; H_1, H_2, H_3 are harmonic functions on (M_0, g^0). The metrics g^i, $i = 0,1,2,3,4,5$, have Euclidean signatures and we put the metric $g^6 = -dt \otimes dt$. The branes sets are $I_1 = \{1,5,6\}$, $I_2 = \{2,4,6\}$, $I_3 = \{3,4,5,6\}$.

The gamma matrices may be chosen in the following form

$$\begin{aligned}(\hat{\Gamma}^A) = (&\hat{\Gamma}^{a_0}_{(0)} \otimes 1 \otimes 1 \otimes \mathbf{1}_2 \otimes 1 \otimes 1 \otimes 1 \otimes \sigma_3 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2, \\ &\mathbf{1}_2 \otimes 1 \otimes 1 \otimes \mathbf{1}_2 \otimes 1 \otimes 1 \otimes 1 \otimes \sigma_1 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2, \\ &\mathbf{1}_2 \otimes 1 \otimes 1 \otimes \mathbf{1}_2 \otimes 1 \otimes 1 \otimes 1 \otimes \sigma_2 \otimes \sigma_3 \otimes \mathbf{1}_2, \\ &\mathbf{1}_2 \otimes 1 \otimes 1 \otimes \hat{\Gamma}^{a_3}_{(3)} \otimes 1 \otimes 1 \otimes 1 \otimes \sigma_2 \otimes \sigma_1 \otimes \mathbf{1}_2, \\ &\mathbf{1}_2 \otimes 1 \otimes 1 \otimes \mathbf{1}_2 \otimes 1 \otimes 1 \otimes 1 \otimes \sigma_2 \otimes \sigma_2 \otimes \sigma_3, \\ &\mathbf{1}_2 \otimes 1 \otimes 1 \otimes \mathbf{1}_2 \otimes 1 \otimes 1 \otimes 1 \otimes \sigma_2 \otimes \sigma_2 \otimes \sigma_1, \\ &\mathbf{1}_2 \otimes 1 \otimes 1 \otimes \mathbf{1}_2 \otimes 1 \otimes 1 \otimes i \otimes \sigma_2 \otimes \sigma_2 \otimes \sigma_2).\end{aligned} \qquad (3.79)$$

Here the operators

$$\hat{\Gamma}_{(0)} = \hat{\Gamma}^{1_0}_{(0)} \hat{\Gamma}^{2_0}_{(0)} \hat{\Gamma}^{3_0}_{(0)}, \quad \hat{\Gamma}_{(3)} = \hat{\Gamma}^{1_3}_{(3)} \hat{\Gamma}^{2_3}_{(3)} \hat{\Gamma}^{3_3}_{(3)} \qquad (3.80)$$

obey

$$(\hat{\Gamma}_{(0)})^2 = (\hat{\Gamma}_{(3)})^2 = -\mathbf{1}_2. \qquad (3.81)$$

The spinor monomial reads

$$\eta = \eta_0(x) \otimes \eta_1(y_1) \otimes \eta_2(y_2) \otimes \eta_3(y_3) \otimes \eta_4(y_4) \otimes \eta_5(y_5) \otimes \eta_6(y_6) \otimes \chi, \qquad (3.82)$$

where $\eta_i = \eta_i(y_i)$ is a 1-component spinor on M_i, $i = 1,2,4,5,6$, $\eta_0 = \eta_0(x)$ is a 2-component spinor on M_0, $\eta_3 = \eta_3(y_3)$ is a 2-component spinor on M_3 and χ belongs to $V = \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$.

The covariant derivatives $\bar{D}^{(i)}_{m_i}$ act on η as follows

$$\begin{aligned}\bar{D}^{(0)}_{m_0} \eta &= \big(D^{(0)}_{m_0} \eta_0\big) \otimes \eta_1 \otimes \eta_2 \otimes \eta_3 \otimes \eta_4 \otimes \eta_5 \otimes \eta_6 \otimes \chi, \\ &\qquad\qquad\qquad\qquad \ldots \\ \bar{D}^{(3)}_{m_3} \eta &= \eta_0 \otimes \eta_1 \otimes \eta_2 \otimes \big(D^{(3)}_{m_3} \eta_3\big) \otimes \eta_4 \otimes \eta_5 \otimes \eta_6 \otimes \chi, \\ &\qquad\qquad\qquad\qquad \ldots \\ \bar{D}^{(6)}_{m_6} \eta &= \eta_0 \otimes \eta_1 \otimes \eta_2 \otimes \eta_3 \otimes \eta_4 \otimes \eta_5 \otimes \big(D^{(6)}_{m_6} \eta_6\big) \otimes \chi,\end{aligned} \qquad (3.83)$$

where

$$\begin{aligned}\bar{D}^{(0)}_{m_0} &= \partial_{m_0} + \frac{1}{4} \omega^{(0)}_{a_0 b_0 m_0} (\hat{\Gamma}^{a_0}_{(0)} \hat{\Gamma}^{b_0}_{(0)} \otimes 1 \otimes 1 \otimes \mathbf{1}_2 \otimes 1 \otimes 1 \otimes 1 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2), \\ \bar{D}^{(3)}_{m_3} &= \partial_{m_3} + \frac{1}{4} \omega^{(3)}_{a_3 b_3 m_3} (\mathbf{1}_2 \otimes 1 \otimes 1 \otimes \hat{\Gamma}^{a_3}_{(3)} \hat{\Gamma}^{b_3}_{(3)} \otimes 1 \otimes 1 \otimes 1 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2),\end{aligned} \qquad (3.84)$$

$\bar{D}^{(i)}_{m_i} = D^{(i)}_{m_i} = \partial_{m_i}$ for $i = 1, 2, 4, 5, 6$ and $D^{(i)}_{m_i}$ are covariant derivatives corresponding to M_i, $i = 0, 3$.

Under (3.79) the projection operators corresponding to the $M2$-branes read

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_1}\hat{\Gamma}^{1_5}\hat{\Gamma}^{1_6} = -\mathbf{1}_2 \otimes 1 \otimes 1 \otimes \mathbf{1}_2 \otimes 1 \otimes 1 \otimes 1 \otimes B_1 \tag{3.85}$$

for $s = I_1$ and

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_2}\hat{\Gamma}^{1_4}\hat{\Gamma}^{1_6} = -\mathbf{1}_2 \otimes 1 \otimes 1 \otimes \mathbf{1}_2 \otimes 1 \otimes 1 \otimes 1 \otimes B_2 \tag{3.86}$$

for $s = I_2$. The operator for the $M5$-brane can be written in the form

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_0}\hat{\Gamma}^{2_0}\hat{\Gamma}^{3_0}\hat{\Gamma}^{1_1}\hat{\Gamma}^{1_2} = i\hat{\Gamma}_{(0)} \otimes 1 \otimes 1 \otimes \mathbf{1}_2 \otimes 1 \otimes 1 \otimes 1 \otimes B_3 \tag{3.87}$$

for $s = I_3$.

In (3.85)-(3.87) B_s are self-adjoint commuting idempotent operators acting on $V = \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$

$$B_1 = \sigma_1 \otimes \mathbf{1}_2 \otimes \sigma_3, \quad B_2 = \sigma_2 \otimes \sigma_3 \otimes \sigma_1, \quad B_3 = \mathbf{1}_2 \otimes \sigma_3 \otimes \mathbf{1}_2. \tag{3.88}$$

The gamma matrices corresponding to M_0 can be chosen in the form $(\hat{\Gamma}^{a_0}_{(0)}) = (\sigma_1, \sigma_2, \sigma_3)$ and hence $\hat{\Gamma}_{(0)} = i\mathbf{1}_2$.

Under the Proposition from Appendix there exists a basis of eigenvectors $\psi_{\varepsilon_1\varepsilon_2\varepsilon_3}$ in V with $\varepsilon_1 = \pm 1$, $\varepsilon_2 = \pm 1$, $\varepsilon_3 = \pm 1$, satisfying (3.72).

The solutions to generalized Killing equations (2.6) corresponding to the field configuration from (3.77), (3.78) are given by the following monomial solutions

$$\varepsilon = H_1^{-1/6} H_2^{-1/6} H_3^{-1/12} \eta_0(x) \otimes \eta_1 \otimes \eta_2 \otimes \eta_3(y_3) \otimes \eta_4 \otimes \eta_5 \otimes \eta_6 \otimes \psi_{\varepsilon_1,\varepsilon_2,\varepsilon_3}, \tag{3.89}$$

where $\eta_0(x)$ and $\eta_3(y_3)$ are parallel spinors defined on M_0 and M_3, respectively, η_i is a constant 1-dimensional spinor on M_i, $i = 1, 2, 4, 5, 6$.

Using the relations (2.20), (3.82) with $\chi = \psi_{\varepsilon_1,\varepsilon_2,\varepsilon_3}$, (3.85)-(3.87) one can obtain the restrictions for the parameters ε_s

$$-\varepsilon_s = c_s. \tag{3.90}$$

Thus the number of linear independent solutions given by (3.89) and (3.90)

$$N = 32\mathcal{N} = n_0 n_3, \tag{3.91}$$

where n_j is the number of parallel spinors on the 3-dimensional manifolds M_j, $j = 0, 3$.

Example.

Let the totally transverse space M_0 and relatively transverse space M_6 be coinciding with 3-dimensional Euclidean space: $M_0 = M_6 = \mathbb{R}^3$. Then one obtains $\mathcal{N} = 1/8$ in agreement with [15].

3.4 $M2 \cap M5 \cap M5$

According to the classification from [15] there are two possible configurations for the intersection of one $M2$-brane and two $M5$-branes.

(i)

The first configuration $M2 \cap M5 \cap M5$ is defined on the manifold

$$M_0 \times M_1 \times M_2 \times M_3 \times M_4 \times M_5, \tag{3.92}$$

where $d_0 = d_2 = d_3 = d_4 = d_5 = 2$ and $d_1 = 1$ and describes a $M2$-brane intersecting each of the two $M5$-branes over a string with the $M5$-branes intersecting over a 3-brane (see Figure 6).



Figure 6: $M2 \cap M5 \cap M5$

The solution for the intersection of an electric $M2$-brane and two magnetic ones is given by

$$g = H_1^{1/3} H_2^{2/3} H_3^{2/3} \Big\{ \hat{g}^0 + H_1^{-1} \hat{g}^1 + H_2^{-1} \hat{g}^2 + H_3^{-1} \hat{g}^3 + H_2^{-1} H_3^{-1} \hat{g}^4 + H_1^{-1} H_2^{-1} H_3^{-1} \hat{g}^5 \Big\}, \tag{3.93}$$

$$F = c_1 dH_1^{-1} \wedge \hat{\tau}_1 \wedge \hat{\tau}_5 + c_2 (*_0 dH_2) \wedge \hat{\tau}_1 \wedge \hat{\tau}_3 + c_3 (*_0 dH_3) \wedge \hat{\tau}_1 \wedge \hat{\tau}_2, \tag{3.94}$$

where $c_1^2 = c_2^2 = c_3^2 = 1$; H_1, H_2, H_3 are harmonic functions defined on (M_0, g^0). The metrics g^i, $i = 0, 1, 2, 3, 4$, have Euclidean signatures and the metric g^5 has the signature $(-,+)$. The branes sets are $I_1 = \{1, 5\}$, $I_2 = \{2, 4, 5\}$ and $I_3 = \{3, 4, 5\}$.

We introduce the following set of Γ-matrices

$$(\hat{\Gamma}^A) = \begin{array}{ccccccccccc} (\hat{\Gamma}^{a_0}_{(0)} & \otimes & 1 & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2, \\ \hat{\Gamma}_{(0)} & \otimes & 1 & \otimes & \hat{\Gamma}_{(2)} & \otimes & \hat{\Gamma}_{(3)} & \otimes & \hat{\Gamma}_{(4)} & \otimes & \hat{\Gamma}_{(5)} \\ i\hat{\Gamma}_{(0)} & \otimes & 1 & \otimes & \hat{\Gamma}^{a_2}_{(2)} & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2 \\ \hat{\Gamma}_{(0)} & \otimes & 1 & \otimes & \hat{\Gamma}_{(2)} & \otimes & \hat{\Gamma}^{a_3}_{(3)} & \otimes & \mathbf{1}_2 & \otimes & \mathbf{1}_2 \\ i\hat{\Gamma}_{(0)} & \otimes & 1 & \otimes & \hat{\Gamma}_{(2)} & \otimes & \hat{\Gamma}_{(3)} & \otimes & \hat{\Gamma}^{a_4}_{(4)} & \otimes & \mathbf{1}_2 \\ \hat{\Gamma}_{(0)} & \otimes & 1 & \otimes & \hat{\Gamma}_{(2)} & \otimes & \hat{\Gamma}_{(3)} & \otimes & \hat{\Gamma}_{(4)} & \otimes & \hat{\Gamma}^{a_5}_{(5)}), \end{array} \tag{3.95}$$

where $\hat{\Gamma}^{a_i}_{(i)}$ are 2×2 Γ-matrices, $a_i = 1_i, 2_i$, $i = 0, 2, 3, 4, 5$, corresponding to M_i, respectively, and the operators

$$\hat{\Gamma}_{(i)} = \hat{\Gamma}^{1_i}_{(i)} \hat{\Gamma}^{2_i}_{(i)}, \quad \hat{\Gamma}_{(5)} = \hat{\Gamma}^{1_5}_{(5)} \hat{\Gamma}^{2_5}_{(5)} \tag{3.96}$$

satisfy

$$(\hat{\Gamma}_{(i)})^2 = -\mathbf{1}_2, \quad (\hat{\Gamma}_{(5)})^2 = \mathbf{1}_2, \tag{3.97}$$

$i = 0, 2, 3, 4$.

Consider η in the form $\eta = \eta_0(x) \otimes \eta_1(y_1) \otimes \eta_2(y_2) \otimes \eta_3(y_3) \otimes \eta_4(y_4) \otimes \eta_5(y_5)$, where $\eta_i = \eta_i(y_i)$ is a 2-component spinor on M_i, $i = 0,2,3,4,5$, $\eta_1(y_1)$ is a 1-component spinor on M_1.

Due to (3.95) and (3.97) the operator $\bar{D}^{(i)}_{m_i}$ acts on η as

$$\bar{D}^{(i)}_{m_i}\eta = \ldots \otimes \eta_{i-1} \otimes \left(D^{(i)}_{m_i}\eta_i\right) \otimes \eta_{i+1} \otimes \ldots, \tag{3.98}$$

where $D^{(i)}_{m_i}$ is the spinorial covariant derivative corresponding to M_i, $i = 0,2,3,4,5$, and $D^{(1)}_{m_1} = \partial_{m_1}$. Thus the relations (2.23) are satisfied if η_i is a parallel spinor on M_i, $i = 0,1,2,3,4,5$, and $\eta_1(y_1) = \eta_1$ is constant.

The operator corresponding $M2$-brane is

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_1}\hat{\Gamma}^{1_5}\hat{\Gamma}^{2_5} = \hat{\Gamma}_{(0)} \otimes 1 \otimes \hat{\Gamma}_{(2)} \otimes \hat{\Gamma}_{(3)} \otimes \hat{\Gamma}_{(4)} \otimes \mathbf{1}_2 \tag{3.99}$$

for $s = I_1$, the operators corresponding to $M5$-branes are

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_0}\hat{\Gamma}^{2_0}\hat{\Gamma}^{1_1}\hat{\Gamma}^{1_3}\hat{\Gamma}^{2_3} = \mathbf{1}_2 \otimes 1 \otimes \hat{\Gamma}_{(2)} \otimes \mathbf{1}_2 \otimes \hat{\Gamma}_{(4)} \otimes \hat{\Gamma}_{(5)} \tag{3.100}$$

for $s = I_2$ and

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_0}\hat{\Gamma}^{2_0}\hat{\Gamma}^{1_1}\hat{\Gamma}^{1_2}\hat{\Gamma}^{2_2} = \mathbf{1}_2 \otimes 1 \otimes \mathbf{1}_2 \otimes \hat{\Gamma}_{(3)} \otimes \hat{\Gamma}_{(4)} \otimes \hat{\Gamma}_{(5)} \tag{3.101}$$

for $s = I_3$.

The projections (2.20) are satisfied if

$$\begin{aligned} \hat{\Gamma}_{(j)}\eta_j = c_{(j)}\eta_j, \qquad c^2_{(j)} = -1, \\ \hat{\Gamma}_{(5)}\eta_5 = c_{(5)}\eta_5, \qquad c^2_{(5)} = 1, \end{aligned} \tag{3.102}$$

$j = 0,3,2,4$, and

$$c_{(0)}c_{(2)}c_{(3)}c_{(4)} = c_1, \quad c_{(2)}c_{(4)}c_{(5)} = c_2, \quad c_{(3)}c_{(4)}c_{(5)} = c_3. \tag{3.103}$$

Equations (3.103) have the following solutions

$$c_{(0)} = ic_1c_2c_3\varepsilon_4, \quad c_{(2)} = -ic_2\varepsilon_4\varepsilon_5, \tag{3.104}$$

$$c_{(3)} = -ic_3\varepsilon_4\varepsilon_5, \quad c_{(4)} = i\varepsilon_4, \quad c_{(5)} = \varepsilon_5, \tag{3.105}$$

where $\varepsilon_4 = \pm 1$, $\varepsilon_5 = \pm 1$.

For the field configuration (3.93) and (3.94) we obtain the following solution to SUSY equations

$$\varepsilon = H_1^{-1/6}H_2^{-1/12}H_3^{-1/12}\eta_0(x) \otimes \eta_1 \otimes \eta_2(y_2) \otimes \eta_2(y_3) \otimes \eta_4(y_4) \otimes \eta_5(y_5), \tag{3.106}$$

where η_i, $i = 0,2,3,4,5$, are chiral parallel spinors defined on M_i $\left(D^{(i)}_{m_i}\eta_i = 0\right)$, obeying (3.102) and (3.103), η_1 is constant.

Thus, the number of linear independent solutions is

$$N = 32\mathcal{N} = \sum_{\substack{\varepsilon_4=\pm1,\\ \varepsilon_5=\pm1}} n_0(i\varepsilon_4 c_1 c_2 c_3) n_2(-ic_2\varepsilon_4\varepsilon_5) n_3(-ic_3\varepsilon_4\varepsilon_5) n_4(i\varepsilon_4) n_5(\varepsilon_5), \qquad (3.107)$$

where $n_j(c_j)$ is the number of chiral parallel spinors on M_j, $j = 0, 2, 3, 4, 5$.

Examples.

Let $M_0 = M_2 = M_3 = M_4 = \mathbb{R}^2$ and $M_1 = \mathbb{R}$. Then for $M_5 = \mathbb{R}^{1,1}$ one gets $\mathcal{N} = 1/8$, while for $M_5 = \mathbb{R}^{1,1}/Z_2$ we find $\mathcal{N} = 1/16$ for any choice of brane sign factors.

(ii)

The second possible intersection of $M2$-brane and two $M5$-branes is defined on the manifold

$$M_0 \times M_1 \times M_2 \times M_3 \times M_4 \times M_5 \times M_6, \qquad (3.108)$$

where $d_0 = 3$, $d_1 = d_2 = d_4 = d_5 = d_6 = 1$, $d_3 = 3$.

The configuration is given in Figure 7.



Figure 7: $M2 \cap M5 \cap M5$

The solution describing the intersection of one electic brane and two magnetic ones reads now

$$g = H_1^{1/3} H_2^{2/3} H_3^{2/3} \Big\{ \hat{g}^0 + H_2^{-1}\hat{g}^1 + H_3^{-1}\hat{g}^2 + H_2^{-1}H_3^{-1}\hat{g}^3 + H_1^{-1}H_3^{-1}\hat{g}^4 + H_1^{-1}H_2^{-1}\hat{g}^5 + H_1^{-1}H_2^{-1}H_3^{-1}\hat{g}^6 \Big\}, \qquad (3.109)$$

$$F = c_1 dH_1^{-1} \wedge \hat{\tau}_4 \wedge \hat{\tau}_5 \wedge \hat{\tau}_6 + c_2(*_0 dH_2) \wedge \hat{\tau}_2 \wedge \hat{\tau}_4 + c_3(*_0 dH_3) \wedge \hat{\tau}_1 \wedge \hat{\tau}_5, \qquad (3.110)$$

where $c_1^2 = c_2^2 = c_3^2 = 1$; H_1, H_2, H_3 are harmonic functions on (M_0, g^0). The metrics g^i, $i = 0, 1, 2, 3, 4, 5$, have Euclidean signatures and we put the metric $g^6 = -dt \otimes dt$. The branes sets are $I_1 = \{4, 5, 6\}$, $I_2 = \{1, 3, 5, 6\}$, $I_3 = \{2, 3, 4, 6\}$.

The corresponding set of gamma matrices matches with (3.79) for two electric and one magnetic branes due to the same space configuration. The expressions for $\hat{\Gamma}_{(i)}$, $(\hat{\Gamma}_{(i)})^2$, $i = 0, 3,$ coincide with (3.80) and (3.81) as well.

The 32-component spinor η can be represented in the form (3.82) $\eta = \eta_0(x) \otimes \eta_1(y_1) \otimes \eta_2(y_2) \otimes \eta_3(y_3) \otimes \eta_4(y_4) \otimes \eta_5 \otimes \eta_6 \otimes \chi$, where $\eta_i = \eta_i(y_i)$ is a 1-component spinor on M_i, $i = 1, 2, 4, 5, 6$, $\eta_0 = \eta_0(x)$ is a 2-component spinor on M_0 and $\eta_3 = \eta_3(y_3)$ is a 2-component spinor on M_3 and χ is an element of $V = \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$.

For covariant derivatives we have relations (3.83) and (3.84). Using the representation of gamma matrices (3.79) the operators corresponding to M-branes are given by

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{14}\hat{\Gamma}^{15}\hat{\Gamma}^{16} = -\mathbf{1}_2 \otimes 1 \otimes 1 \otimes \mathbf{1}_2 \otimes 1 \otimes 1 \otimes 1 \otimes B_1 \qquad (3.111)$$

for $s = I_1$,

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_0}\hat{\Gamma}^{2_0}\hat{\Gamma}^{3_0}\hat{\Gamma}^{1_2}\hat{\Gamma}^{1_4} = i\hat{\Gamma}_0 \otimes 1 \otimes 1 \otimes \mathbf{1}_2 \otimes 1 \otimes 1 \otimes 1 \otimes B_2 \qquad (3.112)$$

for $s = I_2$,

$$\hat{\Gamma}_{[s]} = \hat{\Gamma}^{1_0}\hat{\Gamma}^{2_0}\hat{\Gamma}^{3_0}\hat{\Gamma}^{1_1}\hat{\Gamma}^{1_5} = i\hat{\Gamma}_{(0)} \otimes 1 \otimes 1 \otimes \mathbf{1}_2 \otimes 1 \otimes 1 \otimes 1 \otimes B_3 \qquad (3.113)$$

for $s = I_3$.

Here

$$B_1 = \sigma_2 \otimes \sigma_2 \otimes \mathbf{1}_2, \quad B_2 = \sigma_3 \otimes \sigma_1 \otimes \sigma_3, \quad B_3 = \mathbf{1}_2 \otimes \sigma_2 \otimes \sigma_1 \qquad (3.114)$$

are self-adjoint commuting operators.

As in the case with two $M2$-branes and one $M5$-brane we put $(\hat{\Gamma}^{a_0}_{(0)}) = (\sigma_1, \sigma_2, \sigma_3)$ and hence $\hat{\Gamma}_{(0)} = i\mathbf{1}_2$.

Due to Proposition from Appendix there exists a basis of eigenvectors $\psi_{\varepsilon_1,\varepsilon_2,\varepsilon_3} \in V$ satisfying the relations

$$B_s\psi_{\varepsilon_1,\varepsilon_2,\varepsilon_3} = \varepsilon_s\psi_{\varepsilon_1,\varepsilon_2,\varepsilon_3}, \qquad (3.115)$$

where $\varepsilon_1 = \pm 1$, $\varepsilon_2 = \pm 1$, $\varepsilon_3 = \pm 1$, $s = 1, 2, 3$.

The solutions to eqs. (2.6) corresponding to the field configuration from (3.109), (3.110) are given rise by

$$\varepsilon = H_1^{-1/6}H_2^{-1/12}H_3^{-1/12}\eta_0(x) \otimes \eta_1 \otimes \eta_2 \otimes \eta_3(y_3) \otimes \eta_4 \otimes \eta_5 \otimes \eta_6 \otimes \psi_{\varepsilon_1,\varepsilon_2,\varepsilon_3}, \qquad (3.116)$$

where $\eta_0(x)$ and $\eta_3(y_3)$ are parallel spinors on M_0 and M_3, η_i are constant 1-dimensional spinors, $i = 1, 2, 4, 5, 6$.

The parameters ε_s obey the chirality restrictions

$$-\varepsilon_s = c_s, \qquad (3.117)$$

$s = 1, 2, 3$, following from relations (2.20), (3.65) with $\chi = \psi_{\varepsilon_1,\varepsilon_2,\varepsilon_3}$, (3.111)-(3.113), (3.115)

The number of linear independent solutions given by (3.116)-(3.117) can be computed as follows

$$N = 32\mathcal{N} = n_0 n_3, \qquad (3.118)$$

where n_0 and n_3 are numbers of parallel spinors on the manifolds M_0 and M_3, respectively.

Example.

Here the only example we have is the trivial one: $M_0 = M_2 = \mathbb{R}^3$ with $\mathcal{N} = 1/8$ in agreement with [15].

4 Conclusions

In this paper we have considered the generalized Killing equations in $D = 11$ supergravity for triple M-brane solutions defined on the product of Ricci-flat manifolds. The first configuration with three electric branes $M2 \cap M2 \cap M2$ has been studied earlier in [20], while six others ones: $M2 \cap M2 \cap M5$, $M2 \cap M5 \cap M5$ (two configurations) and $M5 \cap M5 \cap M5$ (three configurations) have been considered here for the first time. Using the approach of [19, 20] we have obtained explicit formulae for computing the amount of preserved supersymmetries for all triple M-brane configurations. These formulae have generalized the relations obtained earlier by several authors for flat factor spaces $\mathbb{R}^{k_i}$ [14, 15]. The deduced fractional numbers of preserved SUSY $\mathcal{N}$ depend upon certain numbers of (chiral or all) parallel spinors on some factor spaces and in several cases upon brane sign factors c_s.

We have presented examples of partially supersymmetric configurations which do not belong to the classification of Bergshoeff et al [15]. These examples use the following factor-spaces: $K3$ [20], $\mathbb{C}^2_*/Z_2$ (for $M2 \cap M2 \cap M2$), $\mathbb{R}^{1,1}_*/Z_2$ (for case (i) of $M2 \cap M5 \cap M5$ and case (ii) of $M5 \cap M5 \cap M5$) and $(\mathbb{R}^{1,1}_*/Z_2) \times \mathbb{R}$ (for case (iii) of $M5 \cap M5 \cap M5$), 4-dimensional pp-wave manifold from [13] and $(\mathbb{R}^{1,1}_*/Z_2) \times \mathbb{R}^2$ (for case (i) of $M5 \cap M5 \cap M5$). This list of factor spaces contains only few 4-dimensional Ricci-flat factor spaces ($K3$, pp-wave) which are not flat. All other two- and three-dimensional factor-spaces are flat. (Any $d = 2, 3$ Ricci-flat space is flat.)

In three cases: $M2 \cap M2 \cap M2$ and $M5 \cap M5 \cap M5$ ((i) and (ii)) we presented examples where $\mathcal{N}$ depend upon brane sign factors $c_s = \pm 1$.

An open problem here is to analyze special solutions with certain "near-horizon" harmonic functions H_s for which the unbroken numbers of supersymmetries might be larger then the numbers $\mathcal{N}$ obtained here for generic H_s-functions. Here we will deal with Freund-Rubin-type solutions with composite M-branes [28]. Such partially supersymmetric solutions will lead to certain relations which contain numbers of (chiral) Killing spinors on certain Einstein factor spaces. This may of interest in a context of AdS/CFT approach, its generalizations and applications [26].

It is also a straightforward task to use the obtained results for studying partially supersymmetric solutions (with Ricci-flat or non-trivial flat factor-spaces) in IIA-, IIB- and other ($d < 10$) supergravitational models using dimensional reductions and duality transformations. Another problem of interest may be related to a search of "pseudo-supersymmetric" brane solutions [27] defined on product of Ricci-flat manifolds by using a possible generalization of the approach from [19, 20].

Acknowledgments

The authors are grateful to D.V. Alekseevsky and H. Baum for helpful comments on related issues. A. G. was supported in part by The Ministry of education and science of Russian Federation, project 14.B37.21.2035.

A Appendix

Here we outline a proposition on simultaneous diagonalization of a set of linear idempotent operators arising in the decomposition of gamma matrices for some cases of product spaces. This proposition is a special case of the so-called " 2^{-k}-splitting" theorem from [19].

Proposition.

Let $B_1, \ldots, B_k : V \to V$ *be a set of linear operators defined on the vector space* $V = \mathbb{C}^2 \otimes ... \otimes \mathbb{C}^2$ *(k-times) which are idempotent*

$$B_i^2 = \mathbf{1}_V \tag{A.1}$$

and commute with each other

$$B_i B_j = B_j B_i, \tag{A.2}$$

$i, j = 1, ..., k$.

There is a set of bijective operators $A_1, \ldots A_k : V \to V$ *obeying the following relations*

$$A_i B_i + B_i A_i = 0, \tag{A.3}$$

$$A_i B_j = B_j A_i, \quad i \neq j, \tag{A.4}$$

$i, j = 1, ..., k$.

Then there exists a basis of 2^k *vectors in* V *,* $\psi_{\varepsilon_1,...,\varepsilon_k}$ *,* $\varepsilon_1, ..., \varepsilon_k = \pm 1$ *, which are eigenvectors of* B_i *:*

$$B_i \psi_{\varepsilon_1,...,\varepsilon_k} = \varepsilon_i \psi_{\varepsilon_1,...,\varepsilon_k}, \tag{A.5}$$

$i = 1, ..., k$.

According to the proof of the theorem in [19], the operator A_i defines an isomorphism between vector eigen-spaces $V_{...,\varepsilon_i,...}$ and $V_{...,-\varepsilon_i,...}$ (all other indices ε_j , $j \neq i$, are coinciding). Hence the basis may be found as follows. First we find a non-zero “ground-state” vector $\psi_{-1,...,-1} = \psi$ satifying

$$B_i \psi = -\psi \tag{A.6}$$

for all i and afterwards we put

$$\psi_{\varepsilon_1,...,\varepsilon_k} \equiv A_1^{(1+\varepsilon_1)/2} \ldots A_k^{(1+\varepsilon_k)/2} \psi, \tag{A.7}$$

where $\varepsilon_1, \ldots, \varepsilon_k = \pm 1$. Here $A_i^0 \equiv \mathbf{1}_V$ is identity operator on V .

Let us consider three examples of the operators B_1, B_2, B_3 , which appeared in the previous section.

Example 1.

Let

$$B_1 = -\sigma_1 \otimes \sigma_2 \otimes \mathbf{1}_2, \quad B_2 = \sigma_3 \otimes \sigma_1 \otimes \sigma_3, \quad B_3 = -\mathbf{1}_2 \otimes \sigma_2 \otimes \sigma_1. \tag{A.8}$$

A set of the operators A_s , $s = 1, 2, 3$, obeying (A.3)-(A.4) may be chosen as

$$A_1 = \sigma_3 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2, \quad A_2 = \sigma_1 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2, \quad A_3 = \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \sigma_3. \tag{A.9}$$

Example 2.

For

$$B_1 = \sigma_1 \otimes \mathbf{1}_2 \otimes \sigma_3, \quad B_2 = \sigma_2 \otimes \sigma_3 \otimes \sigma_1, \quad B_3 = \mathbf{1}_2 \otimes \sigma_3 \otimes \mathbf{1}_2, \tag{A.10}$$

the operators A_s read

$$A_1 = \sigma_2 \otimes \mathbf{1}_2 \otimes \sigma_3, \quad A_2 = \sigma_1 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2, \quad A_3 = \sigma_1 \otimes \sigma_1 \otimes \mathbf{1}_2. \tag{A.11}$$

Example 3.

Let

$$B_1 = \sigma_2 \otimes \sigma_2 \otimes \mathbf{1}_2, \quad B_2 = \sigma_3 \otimes \sigma_2 \otimes \sigma_3, \quad B_3 = \mathbf{1}_2 \otimes \sigma_2 \otimes \sigma_1. \tag{A.12}$$

Then A_s-operators can be presented in the following form

$$A_1 = \sigma_3 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2, \quad A_2 = \sigma_2 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2, \quad A_3 = \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \sigma_3. \tag{A.13}$$

References

[1] J. Bagger and N. Lambert, Modeling multiple M2's, *Phys. Rev.* **D 75**, 045020 (2007); arXiv:hep-th/0611108v3.

[2] A. Gustavsson, Algebraic structures on parallel M2-branes, *Nucl. Phys.* **B 811**, 66 (2009); arXiv:0709.1260v5[hep-th].

[3] O. Aharony, O. Bergman, D. L. Jafferis and J. Maldacena, N=6 superconformal Chern-Simons-matter theories, M2-branes and their gravity duals, *JHEP* **0810**, 091 (2008); arXiv:0806.1218v4 [hep-th].

[4] E. Cremmer, B. Julia and J. Scherk, *Phys. Lett.* **B 76**, 409 (1978).

[5] M. Duff and K. Stelle, Multimembrane solutions of D = 11 supergravity, *Phys. Lett.* **B 253**, 113-118 (1991).

[6] R. Güven, Black p-brane Solutions of D=11 Supergravity Theory, *Phys. Lett.* **B 276**, 49 (1992).

[7] D.J. Smith, Intersecting brane solutions in string and M-theory, topical review, *Class. Quant. Grav.*, **20** R233-317 (2003); arXiv:hep-th/0210157v3.

[8] V.D. Ivashchuk and V.N. Melnikov, Exact solutions in multidimensional gravity with antisymmetric forms, topical review, *Class. Quantum Grav.* **18**, R87-R152 (2001); arXiv:hep-th/0110274.

[9] M.J. Duff, H. Lü, C.N. Pope and E. Sezgin, Supermembranes with Fewer Supersymmetries, *Phys. Lett.* **B 371**, 206 (1996); arXiv:hep-th/9511162.

[10] J.P. Gauntlett, G.W. Gibbons, G. Papadopoulos and P.K. Townsend, Hyper-Kähler manifolds and multiply intersecting branes, *Nucl. Phys.* **B 500**, 133 (1997); arXiv:hep-th/9702202v4.

[11] D. R. Brecher and M.J. Perry, Ricci-Flat Branes, *Nucl. Phys.* B **556**, 151-172 (2000); arXiv:hep-th/9908018.

[12] A. Kaya, New brane solutions from Killing spinor equations, *Nucl. Phys.* **B 583**, 411 (2000); arXiv:hep-th/0004199v2.

[13] J.M. Figueroa-O'Farrill, More Ricci-flat branes, *Phys. Lett.*, B **471** 128-132, (1999); arXiv:hep-th/9910086v2.

[14] A. Tseytlin, Harmonic superpositions of M-branes, *Nucl. Phys.* **B 475**, 149 (1996); arXiv:hep-th/9604035v4.

[15] E. Bergshoeff, M. de Roo, E. Eyras, B. Janssen and J.P. van der Schaar, Multiple Intersections of D-branes and M-branes, *Class. Quantum Grav.*, **14**, 2757 (1997); arXiv:hep-th/9612095v3.

[16] V.D. Ivashchuk and V.N. Melnikov, Sigma-model for the generalized composite p-branes, *Class. Quantum Grav.* **14**, 3001 (1997), Corrigenda ibid. **15**, 3941 (1998); arXiv:hep-th/9705036v2.

[17] K.S. Stelle, Lectures on supergravity p-branes, arXiv:hep-th/9701088.

[18] J.P. Gauntlett, Intersecting branes, *Seoul/Soktcho 1997, Dualities in gauge and string theories* (May, 1997) 146-193, arXiv:hep-th/9705011v2.

[19] V. D. Ivashchuk, On supersymmetric solutions in $D = 11$ supergravity on product of Ricci-flat spaces, *Grav. Cosmol.*, **6**, No 4, 344 (2000); arXiv:hep-th/0012263.

[20] V. D. Ivashchuk, More M-branes on product of Ricci-flat manifolds, *IJGMMP* **9** No 8, 1250067(27p.), (2012); arXiv:1107.4089v3 [hep-th].

[21] M.Y. Wang, Parallel spinors and parallel forms, *Ann. Glob. Anal. and Geom.*, **7**, 59-68 (1989).

[22] H. Baum, Twistor and Killing spinors in Lorentzian geometry, *Seminaires and Congres*, **4**, 35-52 (2000).

[23] D. Alekseevsky, V. Cortes, C. Devchand and U. Semmelmann, Killing Spinors are Killing vector fields in Riemannian supergeometry, *J. Geom. Phys.* **26**, No. 1-2, 51-78 (1998).

[24] C. Bohle, Killing and twistor spinors on Lorentzian manifolds, Diplomarbeit, Freie Universität Berlin, 1998, 133 pp.

[25] H. Lü, C.N. Pope and J. Rahmfeld, A Construction of Killing Spinors on S^n, *J. Math. Phys.* **40**, 4518 (1999); arXiv:hep-th/9805151.

[26] J. Maldacena and D. Martelli, The unwarped, resolved, deformed conifold: fivebranes and the baryonic branch of the Klebanov-Strassler theory, *JHEP* **1001**, 104 (2010); arXiv:0906.0591 [hep-th].

[27] H. Lü and Z.-L. Wang, Pseudo-supersymmetric p-branes, bubbling and less-bubbling AdS spaces, *JHEP* **1106**, 113 (2011); arXiv:1103.0563 [hep-th].

[28] V.D. Ivashchuk, Composite p-branes on product of Einstein spaces, *Phys. Lett.*, **B 434**, No 1, 28-35 (1998); arXiv:hep-th/9704113v3.